

Algebraic properties of some varieties of central loops ^{*†}

Tèmítópé Gbóláhàn Jaiyéolá¹

&

John Olúsolá Adéníran^{2‡}

Abstract

Isotopes of C-loops with unique non-identity squares are shown to be both C-loops and A-loops. The relationship between C-loops and Steiner loops is further studied. Central loops with the weak and cross inverse properties are also investigated. C-loops are found to be Osborn loops if every element in them are squares.

1 Introduction

C-loops are one of the least studied loops. Few publications that have considered C-loops include Fenyves [14], [15], Beg [7], [8], Phillips et. al. [24], [26], [21], [20], Chein [10] and Solarin et. al. [2], [30], [28], [27]. The difficulty in studying them is as a result of the nature of their identities when compared with other Bol-Moufang identities (the element occurring twice on both sides has no other element separating it from itself). Latest publications on the study of C-loops which has attracted fresh interest on the structure include [24], [26], and [21].

LC-loops, RC-loops and C-loops are loops that satisfies the identities $(xx)(yz) = (x(xy))z$, $(zy)(xx) = z((yx)x)$ and $x(y(yz)) = ((xy)y)z$ respectively. Fenyves' work in [15] was completed in [24]. Fenyves proved that LC-loops and RC-loops are defined by three equivalent identities. But in [24] and [25], it was shown that LC-loops and RC-loops are defined by four equivalent identities. Solarin [28] named the fourth identities left middle(LM-) and right middle(RM-) identities and loops that obey them are called LM-loops and RM-loops respectively. These terminologies were also used in [29]. A C-loop is both an LC-loop and an RC-loop ([15]). Their basic properties are found in [26], [15] and [13].

The right and left translation maps on a loop $(L, \cdot)$ are bijections

$R_x : L \rightarrow L$ and $L_x : L \rightarrow L$ defined as $yR_x = y \cdot x$ and $yL_x = x \cdot y$ respectively for all $x, y \in L$.

^{*}2000 Mathematics Subject Classification. Primary 20N05 ; Secondary 08A05

[†]**Keywords :** central loops, central square, weak inverse property, cross inverse property, unique non-identity commutator, associator, square, Osborn loop.

[‡]Corresponding author

Definition 1.1 Let $(L, \cdot)$ be a loop. The left nucleus of L is the set

$$N_\lambda(L, \cdot) = \{a \in L : ax \cdot y = a \cdot xy \ \forall x, y \in L\}.$$

The right nucleus of L is the set

$$N_\rho(L, \cdot) = \{a \in L : y \cdot xa = yx \cdot a \ \forall x, y \in L\}.$$

The middle nucleus of L is the set

$$N_\mu(L, \cdot) = \{a \in L : ya \cdot x = y \cdot ax \ \forall x, y \in L\}.$$

The nucleus of L is the set

$$N(L, \cdot) = N_\lambda(L, \cdot) \cap N_\rho(L, \cdot) \cap N_\mu(L, \cdot).$$

The centrum of L is the set

$$C(L, \cdot) = \{a \in L : ax = xa \ \forall x \in L\}.$$

The center of L is the set

$$Z(L, \cdot) = N(L, \cdot) \cap C(L, \cdot).$$

L is said to be a centrum square loop if $x^2 \in C(L, \cdot)$ for all $x \in L$. L is said to be a central square loop if $x^2 \in Z(L, \cdot)$ for all $x \in L$. L is said to be left alternative if for all $x, y \in L$, $x \cdot xy = x^2y$ and is said to right alternative if for all $x, y \in L$, $yx \cdot x = yx^2$. Thus, L is said to be alternative if it is both left and right alternative. The triple (U, V, W) such that $U, V, W \in \text{SYM}(L, \cdot)$ is called an autotopism of L if and only if

$$xU \cdot yV = (x \cdot y)W \ \forall x, y \in L.$$

$\text{SYM}(L, \cdot)$ is called the permutation group of the loop $(L, \cdot)$. The group of autotopisms of L is denoted by $\text{AUT}(L, \cdot)$. Let $(L, \cdot)$ and $(G, \circ)$ be two distinct loops. The triple $(U, V, W) : (L, \cdot) \rightarrow (G, \circ)$ such that $U, V, W : L \rightarrow G$ are bijections is called a loop isotopism if and only if

$$xU \circ yV = (x \cdot y)W \ \forall x, y \in L.$$

We investigate central loops with the unique non-identity commutators, associators and squares and those with unique non-trivial squares are also found to be C-loops whose isotopes are C-loops and A-loops. The relationship between C-loops and Steiner loops is further studied.

Central loops with the weak and cross inverse properties are also investigated. C-loops are found to be Osborn loops if every element in them are squares, hence the possibility for a non-associative Osborn C-loop to exist is envisaged.

For definition of concepts in theory of loops readers may consult [9], [29] and [23].

2 Preliminaries

Definition 2.1 ([16]) Let a, b and c be three elements of a loop L . The loop commutator of a and b is the unique element (a, b) of L which satisfies $ab = (ba)(a, b)$ and the loop associator of a, b and c is the unique element (a, b, c) of L which satisfies $(ab)c = \{a(bc)\}(a, b, c)$.

If X, Y , and Z are subsets of a loop L , we denote by (X, Y) and (X, Y, Z) , respectively, the set of all commutators of the form (x, y) and all the associators of the form (x, y, z) , where $x \in X, y \in Y, z \in Z$.

Definition 2.2 ([16]) A 'unique non-identity commutator' is an element $s \neq e$ (e is the identity element) in a loop L with the property that every commutator in L is e or s .

A 'unique non-identity commutator associator' is an element $s \neq e$ in a loop L with the property that every commutator in L is e or s and every associator is e or s .

A 'unique non-identity square' or 'non-trivial square' is an element $s \neq e$ in a loop L with the property that every square in L is e or s .

Definition 2.3 A loop $(L, \cdot)$ is called a weak inverse property loop (W. I. P. L.) if and only if it obeys the weak inverse property (W. I. P.) : $y(xy)^\rho = x^\rho$ for all $x, y \in L$ while L is called a cross inverse property loop (C. I. P. L.) if and only if it obeys the cross inverse property (C. I. P.) : $xy \cdot x^\rho = y$. $(L, \cdot)$ is called a left inverse property loop or right inverse property loop (L.I.P.L. or R.I.P.L.) if and only if it obeys the left inverse property or right inverse property (L.I.P. or R.I.P.): $x^\lambda(xy) = y$ or $(yx)x^\rho = y$. Hence, it is called an inverse property loop (I.P.L.) if and only if it has the inverse property (I.P.) i.e. it is both an L.I.P. and an R.I.P. loop.

Most of our results and proofs, are stated and written in dual form relative to RC-loops and LC-loops. That is, a statement like 'LC(RC)-loop... A(B)' where 'A' and 'B' are some equations or expressions simply means 'A' is for LC-loops while 'B' is for RC-loops. This is done so that results on LC-loops and RC-loops can be combined to derive those on C-loops. For instance an LC(RC)-loop is a L.I.P.L.(R.I.P.L) loop while a C-loop is an I.P.L. loop.

3 Inner mappings

Lemma 3.1 Let L be a C-loop. Then for each $(A, B, C) \in \text{AUT}(L, \cdot)$, there exists a unique pair of $(S_1, T_1, R_1), (S_2, T_2, R_2) \in \text{AUT}(L, \cdot)$ for each $x \in L$ such that $L_x^2 = S_2^{-1}S_1, R_x^2 = T_1^{-1}T_2, R_x^{-2}L_x^2 = R_2^{-1}R_1, R_1^{-1}R_2T_2^{-1}T_1S_2^{-1}S_1 = I$.

Proof

If L is a C-loop, then $(L_x^2, I, L_x^2), (I, R_x^2, R_x^2) \in \text{AUT}(L)$ for all $x \in L$. So there exist $(S_1, T_1, R_1), (S_2, T_2, R_2) \in \text{AUT}(L, \cdot)$ such that

$$(S_1, T_1, R_1) = (A, B, C)(L_x^2, I, L_x^2) \in \text{AUT}(L)$$

$$(S_2, T_2, R_2) = (A, B, C)(I, R_x^2, R_x^2) \in \text{AUT}(L).$$

Hence, the conditions hold although the identities do not depend on (A, B, C) , but the uniqueness does. ♠

Theorem 3.1 *Let L be a C-loop and let there exist a unique pair of autotopisms $(S_1, T_1, R_1), (S_2, T_2, R_2)$ such that the conditions $L_x^2 = S_2^{-1}S_1, R_x^2 = T_1^{-1}T_2$ and $R_x^{-2}L_x^2 = R_2^{-1}R_1$ hold for each fixed $x \in L$. If $\alpha_1 = S_1^{-1}, \alpha_2 = S_2^{-1}, \beta_1 = T_1^{-1}, \beta_2 = T_2^{-1}, \gamma_1 = R_1^{-1}$ and $\gamma_2 = R_2^{-1}$, then:*

$$(x^2y)\alpha_1 = y\alpha_2, \quad (yx^2)\beta_2 = y\beta_1, \quad (x^2yx^{-2})\gamma_1 = y\gamma_2 \quad \forall x, y \in L.$$

Proof

From Lemma 3.1:

$$L_x^2 = S_2^{-1}S_1, R_x^2 = T_1^{-1}T_2, R_x^{-2}L_x^2 = R_2^{-1}R_1.$$

Keeping in mind that a C-loop is power associative and nuclear square, we have the following proofs.

1. $L_x^2 = S_2^{-1}S_1$ implies $yL_x^2 = yS_2^{-1}S_1$ for all $y \in L$ implies $yL_{x^2} = yS_2^{-1}S_1$ implies $x^2y = yS_2^{-1}S_1$ implies $(x^2y)S_1^{-1} = yS_2^{-1}$ implies $x^2y\alpha_1 = y\alpha_2$.
2. $R_x^2 = T_1^{-1}T_2$ implies $yR_x^2 = yT_1^{-1}T_2$ for all $y \in L$ implies $yx^2 = yT_1^{-1}T_2$ implies $yx^2T_2^{-1} = yT_1^{-1}$ implies $yx^2\beta_2 = y\beta_1$.
3. $R_x^{-2}L_x^2 = R_2^{-1}R_1$ implies $yR_x^{-2}L_x^2 = yR_2^{-1}R_1$ for all $y \in L$ implies $x^2yx^{-2} = yR_2^{-1}R_1$ implies $(x^2yx^{-2})R_1^{-1} = yR_2^{-1}$ implies $(x^2yx^{-2})\gamma_1 = y\gamma_2$.

♠

Corollary 3.1 *Let L be a C-loop. An autotopism of L can be constructed if there exists at least an $x \in L$ such that $x^2 \neq e$. The inverse can also be constructed.*

Proof

We need Lemma 3.1 and Theorem 3.1. If $x^2 = e$, then the autotopism is trivial. Since L is a C-loop, using Lemma 3.1 and Theorem 3.1, it will be noticed that $(\alpha_1S_2, \beta_1T_2, \gamma_1R_2) \in \text{AUT}(L)$ and $(\alpha_2S_1, \beta_2T_1, \gamma_2R_1) = (\alpha_1S_2, \beta_1T_2, \gamma_1R_2)^{-1}$. Hence the proof. ♠

Lemma 3.2 *Let $(L, \cdot)$ be a C-loop. Then the third component mapping $\gamma_2R_1 : L \rightarrow L$ defined by $y\gamma_2R_1 = x^2yx^{-2}$ for all $x \in L$ of the constructed autotopism $(\alpha_2S_1, \beta_2T_1, \gamma_2R_1) \in \text{AUT}(L)$ is :*

1. an automorphism,
2. a semi-automorphism,
3. a middle inner mapping,

4. a pseudo-automorphism with companion x^2 .

Proof

1. $\gamma_2 R_1$ is already a bijection by the construction of the autotopism $(\alpha_2 S_1, \beta_2 T_1, \gamma_2 R_1) \in \text{AUT}(L)$. So we need only to show that it is an homomorphism. Let $y_1, y_2 \in L$, then : $(y_1 y_2) \gamma_2 R_1 = (x^2 y_1 x^{-2})(x^2 y_2 x^{-2}) = y_1 \gamma_2 R_1 \cdot y_2 \gamma_2 R_1$. Whence, $\gamma_2 R_1$ is an automorphism.
2. It will be observed that, $e \gamma_1 = e \gamma_2$, hence $e \gamma_2 R_1 = e$. $(zy \cdot z) \gamma_2 R_1 = x^2(zy \cdot z)x^{-2} = x^2((zy \cdot z)x^{-2}) = (x^2 z x^{-2})(x^2 y x^{-2}) \cdot z \gamma_2 R_1 = (z \gamma_2 R_1 \cdot y \gamma_2 R_1) \cdot z \gamma_2 R_1$. Thence, $\gamma_2 R_1$ is a semi-automorphism
3. We have already settled the fact that $e \gamma_2 R_1 = e$. $y \gamma_2 R_1 = y R_{x^{-2}} L_{(x^{-2})^{-1}} = y T(x^{-2})$ for all $y \in L$ implies $\gamma_2 R_1 = T(x^{-2}) \in \text{Inn}(L)$. Hence $\gamma_2 R_1$ is a middle inner mapping.
4. This is true from (1) by using the fact that says an automorphism in a C-loop L is a pseudo-automorphism with companion x^2 for all $x \in L$.



Lemma 3.3 *Let $(L, \cdot)$ be a C-loop. Then :*

1. $T(x^{-1}) = R_x T(x^{-2}) L_x^{-1}, T(x)^2 = R_x T(x^{-1})^{-1} L_x^{-1},$
2. $T(x^n) = R_x^{n-1} T(x) L_x^{1-n}, T(x^{-n}) = R_x^{1-n} T(x^{-1}) L_x^{n-1}$ for all $n \in \mathbf{Z}^+,$
3. $R(x, x) = I, L(x, x) = I.$

Proof

1. Using the mapping $\gamma_2 R_1$ in Lemma 3.2, defined by $y \gamma_2 R_1 = x^2 y x^{-2}$ for all $x, y \in L$, $y \gamma_2 R_1 = x^2 y x^{-2} = y R_{x^{-2}} L_{x^2} = y R_x^{-1} R_x^{-1} L_x L_x = y R_x^{-1} T(x^{-1}) L_x$. Thus, $\gamma_2 R_1 = R_x^{-1} T(x^{-1}) L_x$. In Lemma 3.2, we showed that $\gamma_2 R_1$ is the middle inner mapping $T(x^{-2})$. Hence, $T(x^{-2}) = R_x^{-1} T(x^{-1}) L_x$ implies $T(x^{-1}) = R_x T(x^{-2}) L_x^{-1}$. $T(x)^2 = R_x L_x^{-1} R_x L_x^{-1} = R_x (R_{x^{-1}} L_{x^{-1}}^{-1})^{-1} L_x^{-1} = R_x T(x^{-1})^{-1} L_x^{-1}$.
2. We prove these by induction. When

$$n = 1, T(x) = R_x^{1-1} T(x) L_x^{1-1} = R_{x^0} T(x) L_{x^0} = T(x) \text{ for all } x \in L,$$

$$n = 2, T(x^2) = T(xx) = R_{x^2} L_{x^2}^{-1} = R_x R_x L_x^{-1} L_x^{-1} = R_x T(x) L_x^{-1} \text{ for all } x \in L,$$

$$\begin{aligned} n = 3, T(x^3) &= T(x^2 x) = R_{x^2 x} L_{(x^2 x)^{-1}} = R_{x^2} R_x L_{x^{-1} x^{-2}} = R_{x^2} R_x L_{x^{-1}} L_{x^{-2}} \\ &= R_x^2 T(x) L_x^{-2} \text{ for all } x \in L. \end{aligned}$$

Let $n = k$, $T(x^k) = R_x^{k-1}T(x)L_x^{1-k}$.

If $n = k + 1$, $T(x^{k+1}) = T(x^{k-1}x^2) = R_{x^{k-1}x^2}L_{(x^{k-1}x^2)}^{-1} = R_{x^{k-1}x^2}L_{x^{-2}x^{1-k}} = R_{x^{k-1}}R_{x^2}L_{x^{-2}}L_{x^{1-k}} = R_{x^{k-1}}T(x^2)L_{x^{1-k}} = R_x^{k-1}R_xT(x)L_x^{-1}L_x^{1-k} = R_x^kT(x)L_x^{-k}$. Thus, $T(x^n) = R_x^{n-1}T(x)L_x^{1-n}$ for all $n \in \mathbf{Z}^+$. Replace x by x^{-1} , then $T(x^{-n}) = T((x^{-1})^n) = R_{x^{-1}}^{n-1}T(x^{-1})L_{x^{-1}}^{1-n} = R_x^{1-n}T(x^{-1})L_x^{n-1}$. Thus, $T(x^{-n}) = R_x^{1-n}T(x^{-1})L_x^{n-1}$ for all $n \in \mathbf{Z}^+$.

3. The right and left inner mappings as defined gives : $R(x, x) = R_x^2R_x^{-2} = I, L(x, x) = L_x^2L_x^{-2} = I$.



Remark 3.1 Lemma 3.2 gives an example of a bijective mapping that is an automorphism, pseudo-automorphism, semi-automorphism and an inner mapping.

4 Relationship between C-loops and Steiner loops

For a loop $(L, \cdot)$, the bijection $J : L \rightarrow L$ is defined by $xJ = x^{-1}$ for all $x \in L$. L is called a Steiner loop if and only if

$$x^2 = e, \quad yx \cdot x = y \text{ and } xy = yx \quad \forall x, y \in L.$$

Theorem 4.1 In a C-loop $(L, \cdot)$, if any of the following is true for all $z \in L$:

1. $(I, L_z^2, JL_z^2J) \in AUT(L)$,
2. $(R_z^2, I, JR_z^2J) \in AUT(L)$,

then, L is a loop of exponent 4.

Proof

1. If $(I, L_z^2, JL_z^2J) \in AUT(L)$ for all $z \in L$, then : $x \cdot yL_z^2 = (xy)JL_z^2J$ for all $x, y, z \in L$ implies $x \cdot z^2y = xy \cdot z^{-2}$ implies $z^2y \cdot z^2 = y$. Then $y^4 = e$. Hence L is a C-loop of exponent 4.
2. If $(R_z^2, I, JR_z^2J) \in AUT(L)$ for all $z \in L$, then : $xR_z^2 \cdot y = (xy)JR_z^2J$ for all $x, y, z \in L$ implies $(xz^2) \cdot y = [(xy)^{-1}z^2]^{-1}$ implies $(xz^2) \cdot y = z^{-2}(xy)$ implies $(xz^2) \cdot y = z^{-2}x \cdot y$ implies $xz^2 = z^{-2}x$ implies $z^4 = e$. Hence L is a C-loop of exponent 4.



Theorem 4.2 In a C-loop L , if the following are true for all $z \in L$:

1. $(I, L_z^2, JL_z^2J) \in AUT(L)$,

$$2. (R_z^2, I, JR_z^2 J) \in AUT(L),$$

then, L is a central square C -loop of exponent 4.

Proof

By the first hypothesis, If $(I, L_z^2, JL_z^2 J) \in AUT(L)$ for all $z \in L$, then : $x \cdot yL_z^2 = (xy)JL_z^2 J$ for all $x, y, z \in L$ implies $x \cdot z^2 y = xy \cdot z^{-2}$.

By the second hypothesis, If $(R_z^2, I, JR_z^2 J) \in AUT(L)$ for all $z \in L$, then : $xR_z^2 \cdot y = (xy)JR_z^2 J$ for all $x, y, z \in L$ implies $xz^2 \cdot y = z^{-2}(xy)$.

Using the two results above and keeping in mind that L is a C -loop we have :

$x \cdot z^2 y = xz^2 \cdot y$ if and only if $xy \cdot z^{-2} = z^{-2} \cdot xy$. Let $t = xy$ then $tz^{-2} = z^{-2}t$ if and only if $z^2 t^{-1} = t^{-1} z^2$. Let $s = t^{-1}$ then $z^2 \in C(L, \cdot)$ for all $z \in L$.

Since s is arbitrary in L , then the last result shows that L is centrum square. Furthermore, C -loops have been found to be nuclear square in [26], thus $z^2 \in Z(L, \cdot)$. Hence L is a central square C -loop. Finally, by Theorem 4.1, $x^4 = e$. ♠

Corollary 4.1 *In a C -loop $(L, \cdot)$, if $(I, L_z^2, JL_z^2 J) \in AUT(L)$, and $(R_z^2, I, JR_z^2 J) \in AUT(L)$ for all $z \in L$, then the following are true :*

1. L is flexible.
2. $(xy)^2 = (yx)^2$ for all $x, y \in L$.
3. $x \mapsto x^3$ is an anti-automorphism.

Proof

This follows by Theorem 4.2, Lemma 5.1 and Corollary 5.2 of [21]. ♠

Theorem 4.3 *A central square C -loop of exponent 4 is a group.*

Proof

To prove this, it shall be shown that the right inner mapping

$R(x, y) = I$ for all $x, y \in L$. Corollary 4.1 is used. Let $w \in L$.

$wR(x, y) = wR_x R_y R_{xy}^{-1} = (wx)y \cdot (xy)^{-1} = (wx)(x^2 y x^2) \cdot (xy)^{-1} = (wx^3)(yx^2) \cdot (xy)^{-1} = (w^2(w^3 x^3))(yx^2) \cdot (xy)^{-1} = (w^2(xw)^3)(yx^2) \cdot (xy)^{-1} = w^2(xw)^3 \cdot (yx^2)(xy)^{-1} = w^2(xw)^3 \cdot [y \cdot x^2(xy)^{-1}] = w^2(xw)^3 \cdot [y \cdot x^2(y^{-1}x^{-1})] = w^2(xw)^3 \cdot [y(y^{-1}x^{-1} \cdot x^2)] = w^2(xw)^3 \cdot [y(y^{-1}x)] = w^2(xw)^3 \cdot x = w^2(w^3 x^3) \cdot x = w^2 \cdot (w^3 x^3)x = w^2 \cdot (w^3 x^{-1})x = w^2 w^3 = w^5 = w$ if and only if $R(x, y) = I$ if and only if $R_x R_y R_{xy}^{-1} = I$ if and only if $R_x R_y = R_{xy}$ if and only if $zR_x R_y = zR_{xy}$ if and only if $zx \cdot y = z \cdot xy$ if and only if L is a group. Hence the claim is true. ♠

Corollary 4.2 *In a C -loop $(L, \cdot)$, if $(I, L_z^2, JL_z^2 J) \in AUT(L)$, and $(R_z^2, I, JR_z^2 J) \in AUT(L)$ for all $z \in L$, then L is a group.*

Proof

This follows from Theorem 4.2 and Theorem 4.3. ♠

Remark 4.1 *Central square C-loops of exponent 4 are A-loops.*

Theorem 4.4 *Let L be a C-loop. L is a central square loop if and only if $\gamma_2 R_1 = I$.*

Proof

Let $\gamma_2 R_1 = I$ if and only if $T(x^{-2}) = I$ for all $x \in L$ if and only if $R_{x^{-2}} L_{x^2} = I$ if and only if $yx^2 = x^2y$ if and only if L is central square. ♠

Theorem 4.5 *Let L be a C-loop such that the mapping $x \mapsto T(x)$ is a bijection, then L is of exponent 2 if and only if $\gamma_2 R_1 = I$.*

Proof

Let $\gamma_2 R_1 = I$ if and only if $T(x^{-2}) = I$ for all $x \in L$ if and only if $T(x^{-2}) = I = R_x^{-1} T(x^{-1}) L_x$ by Lemma 3.3 if and only if $T(x^{-1}) = T(x)$ if and only if $x^{-1} = x$ since $x \mapsto T(x)$ is a bijection if and only if L is a loop of exponent 2. ♠

Corollary 4.3 *A C-loop in which $x \mapsto T(x)$ is a bijection is of exponent 2 if and only if it is central square.*

Proof

This follows by Theorem 4.4 and Theorem 4.5. ♠

Remark 4.2 *This Corollary leads to answer one of the questions we wish to answer in this work. This we give below.*

Corollary 4.4 *In a central square C-loop L , the map $x \mapsto T(x)$ is a bijection implies L is a Steiner loop.*

Proof

By the converse of Corollary 4.3, a C-loop in which $x \mapsto T(x)$ is a bijection, is of exponent 2 if it is central square. By the result in [26], that an inverse property loop of exponent 2 is exactly a Steiner loop and the fact that C-loops are inverse property loops([26]), L is a Steiner loop. ♠

Corollary 4.5 *In a C-loop $(L, \cdot)$, if*

1. $x \mapsto T(x)$ is a bijection,
2. $(I, L_z^2, J L_z^2 J) \in \text{AUT}(L)$ and
3. $(R_z^2, I, J R_z^2 J) \in \text{AUT}(L)$ for all $z \in L$,

then L is a Steiner loop of exponent 4.

Proof

According to Theorem 4.2, L is a central square loop. Hence, since by hypothesis, $x \mapsto T(x)$ is a bijection, then by Corollary 4.4, L is a Steiner loop. By Theorem 4.1, it has an exponent of 4. ♠

Corollary 4.6 *Let $(L, \cdot)$ be a C-loop such that the mapping $x \mapsto T(x)$ is a bijection. L is a Steiner loop if and only if L is a central square C-loop.*

Proof

It was shown in [26] that Steiner loops are C-loops. Recall that Steiner loops are loops of exponent two by definition, hence by Corollary 4.3, L is central square since in L , the mapping $x \mapsto T(x)$ is a bijection. Conversely : By Corollary 4.3, a central square C-loop L in which the mapping $x \mapsto T(x)$ is a bijection is of exponent two. Whence using the fact in [26] that an inverse property loop of exponent two is a steiner loop, the proof is complete.

♠

4.1 Flexibility in C-loops

Condition(s) under which a C-loop is flexible is given below.

Lemma 4.1 *A C-loop is flexible if the mapping $x \mapsto x^2$ is onto.*

Proof

Let L be a C-loop . Then $yx^2 \cdot y = y \cdot x^2y$ for all $x, y \in L$. Thus, L is square flexible, hence by the observation in [12] , L is flexible since the mapping $x \mapsto x^2$ is onto.

Theorem 4.6 *A C-loop L is flexible if $(I, L_z^2, JL_z^2J), (R_z^2, I, JR_z^2J) \in \text{AUT}(L)$ for all $z \in L$ and the middle inner mappings are of order 2.*

Proof

By Lemma 3.3,

$T(x)^2 = R_x T(x^{-1})^{-1} L_x^{-1} = R_x (R_x T(x^{-2}) L_x^{-1})^{-1} L_x^{-1} = R_x (L_x (R_x T(x^{-2}))^{-1}) L_x^{-1} = R_x (L_x T(x^{-2})^{-1} R_x^{-1}) L_x^{-1} = R_x L_x T(x^{-2})^{-1} R_x^{-1} L_x^{-1} = R_x L_x T(x^{-2})^{-1} (L_x R_x)^{-1}$. So $T(x)^2 = R_x L_x T(x^{-2})^{-1} (L_x R_x)^{-1}$ if and only if $T(x)^2 L_x R_x = R_x L_x T(x^{-2})^{-1} = R_x L_x (\gamma_2 R_1)^{-1} = R_x L_x \gamma_1 R_2$ if and only if $T(x)^2 L_x R_x = R_x L_x \gamma_1 R_2$. If $|T(x)| = 2, T(x)^2 = I$ and if $\gamma_1 R_2 = I$ if and only if L is central square then $L_x R_x = R_x L_x$ if and only if $xy \cdot x = x \cdot yx$ is a flexible loop. ♠

Philips and Vojtěchovský [26] studied the close relationship between C-loops and Steiner loops. In [23], it is shown that Steiner loops are exactly commutative inverse property loops of exponent 2. But in [26], this fact was improved, so that commutativity is not a sufficient condition for an inverse property loop of exponent 2 to be a Steiner loop. So they said ‘Steiner loops are exactly inverse property loops of exponent 2’. This result is general for inverse property loops among which are C-loops. They also proved that Steiner loops are C-loops. Corollary 4.4 give conditions (apart from exponent of 2) under which a C-loop is a Steiner loop while Corollary 4.6 give necessary and sufficient conditions for C-loops to be Steiner loops.

It has been found out in [23] that flexibility is possible in a C-loop if the loop is commutative or diassociative. But C-loops naturally do not even satisfy the latter. Apart from

the condition stated in Lemma 4.1, Theorem 4.6 when compared with Corollary 5.2 of [21] shows that some middle inner-mappings do not need to be of exponent of 2 for a C-loop to be flexible.

5 Unique non-identity Commutator Associator Square

Lemma 5.1 *If s is a unique non-identity commutator in a C-loop L , $|s| = 2$, $s \in C(L)$ and $s \in Z(L^2)$.*

Proof

$xy = (yx)(x, y) \Leftrightarrow (x, y) = (yx)^{-1}(xy) = (x^{-1}y^{-1})(xy)$. $(x, y)^{-1} = [(x^{-1}y^{-1})(xy)]^{-1} = (xy)^{-1}(x^{-1}y^{-1})^{-1} = (y^{-1}x^{-1})(yx) = (y, x)$. Thus, $s^{-1} = s$ or $s^{-1} = e$ implies $s^2 = e$ or $s = e$ implies $s^2 = e$.

If $xs \neq sx$, then $xs = (sx)s$ implies $x = sx$ implies $s = e$ implies $xs = sx$ implies $s \in C(L)$. Hence, $s \in Z(L^2)$. ♠

Lemma 5.2 *If s is a unique non-identity associator in a C-loop L , $s \in N(L)$.*

Proof

If $(xy)s \neq x(ys)$, then $(xy)s = x(ys) \cdot s$ implies $xy = x \cdot ys$ implies $y = ys$ implies $s = e$ implies $(xy)s = x(ys)$ implies $s \in N(L)$. ♠

Lemma 5.3 *Let $(L, \cdot)$ be a C-loop which has a unique non-identity commutator associator s . Then s is a central element of order 2.*

Proof

We shall keep in mind that L as a C-loop has the inverse property. $s \in (L, L)$ implies $s^{-1} \in (L, L)$ implies $s^{-1} = s$ since $s^{-1} \neq e$ if and only if $s \neq e$, hence $s^2 = e$. Let $xs \neq sx$ for some $x, y \in L$, then $xs = (sx)s$ implies $x = sx$ implies $s = e$, which is a contradiction. Thus, $s \in C(L)$. If $(xy)s \neq x(ys)$ for some $x, y \in L$, then $(xy)s = (x \cdot ys)s$ implies $xy = x \cdot ys$ implies $y = ys$ implies $s = e$, which is a contradiction. Thus $s \in N(L)$. Therefore $s \in C(L)$, $s \in N(L)$ implies $s \in Z(L)$. ♠

Remark 5.1 *The result in Lemma 5.3 is similar to that proved for Moufang loops in [16].*

Lemma 5.4 *In an LC-loop or RC-loop $(L, \cdot)$ with a unique non-identity square s , the following are true.*

1. $|s| = 2$.
2. $|x| = 4$ or $|x| = 2$.
3. $s \in N_\lambda$ or $s \in N_\rho$ and $s \in N_\mu$.

Proof

By hypothesis, $x^2 = s$ for all $x \in L$.

1. Since $s \in L$, $s^2 = s$ implies $s^{-1}s^2 = s^{-1}s$ or $s^2s^{-1} = ss^{-1}$ implies $s = e$. This is a contradiction, thus $s^2 = e$ if and only if $|s| = 2$.
2. $x^2 = s$ implies $x^4 = x^2x^2 = s^2 = e$ implies $x^4 = e$ or $x^2 = e$.
3. In an LC-loop, $x^2 \in N_\lambda, N_\mu$, thus $s \in N_\lambda, N_\mu$. In an RC-loop, $x^2 \in N_\rho, N_\mu$, thus $s \in N_\rho, N_\mu$.



Lemma 5.5 *Let $(L, \cdot)$ be a loop.*

1. *If L is an LC-loop, L has a unique non-identity square s if and only if $J = R_s^{-1} = R_{s^{-1}}^{-1}$ or $J = I$.*
2. *If L is an RC-loop, L has a unique non-identity square s if and only if $J = L_s^{-1} = L_{s^{-1}}^{-1}$ or $J = I$.*

Proof

1. $x^2 = s$ if and only if $x^{-1}x^2 = x^{-1}s$ if and only if $x = x^{-1}s = xJR_s$ if and only if $JR_s = I$ if and only if $J = R_s = R_{s^{-1}}^{-1}$.
 $x^2 = e$ if and only if $x = x^{-1}$ if and only if $x = xJ$ if and only if $J = I$.
2. $x^2 = s$ if and only if $x^2x^{-1} = sx^{-1}$ if and only if $x = sx^{-1}$ if and only if $x = xJL_s$ if and only if $I = JL_s$ if and only if $J = L_s^{-1} = L_{s^{-1}}^{-1}$.
 $x^2 = e$ if and only if $x = x^{-1}$ if and only if $x = xJ$ if and only if $J = I$.



Theorem 5.1 *Let $(L, \cdot)$ be a L. I. P. (R. I. P.) RC(LC)-loop with a unique non-identity square s . The following are true.*

1. $s \in Z(L, \cdot)$, hence L is centrum square.
2. $J = L_s(J = R_s)$.
3. $x^2y^2 \neq (xy)^2 \neq y^2x^2 : x \mapsto x^2$ is neither an automorphism nor an anti-automorphism.
4. $(a, b, c) = (bc \cdot a)(ab \cdot c) :$
 (a) $ab = a^{-1}b^{-1}$ if and only if $(J, J, I) \in AUT(L, \cdot)$.
 (b) $(a, b, a) = (bs)(ab \cdot a)$ or $(a, b, a) = b(ab \cdot a)$.
5. L is a group or Steiner loop.

6. If L is not commutative, s is the unique non-identity commutator of L for the case of been a C -loop.

Proof

1. $x^2 = s$ implies $x = sx^{-1}$ implies $x^{-1} = s^{-1}x$. Using the result in Lemma 2.1 of [1], $x^{-1} = (sx^{-1})^{-1} = (x^{-1})^{-1}s^{-1}$ implies $x^{-1} = xs^{-1}$. Thus, $x^{-1} = s^{-1}x = xs^{-1}$ implies $sx = xs$ implies $s \in Z(L, \cdot)$.
2. This follows from Lemma 5.5.
3. If $(xy)^2 = x^2y^2$ or $(xy)^2 = y^2x^2$ then $s = s^2$ implies $s = e$ which is a contradiction.
4. $(a, b, c) = [a(bc)]^{-1} \cdot (ab)c = (bc)^{-1}a^{-1} \cdot (ab)c = (c^{-1}b^{-1})a^{-1} \cdot (ab \cdot c) = [s^{-1}(bc)](s^{-1}a) \cdot (ab \cdot c) = (bc \cdot s^{-1})(s^{-1}a) \cdot (ab \cdot c) = (bcs^{-2} \cdot a)(ab \cdot c) = (bc \cdot a)(ab \cdot c)$. So, $(a, b, c) = (bc \cdot a)(ab \cdot c)$.
 (a) Put $c = e$, $(a, b, e) = (ba)(ab) = e$ if and only if $(ab) = (ba)^{-1}$ if and only if $ab = a^{-1}b^{-1}$ if and only if $(J, J, I) \in AUT(L, \cdot)$.
 (b) Let $c = a$, $(a, b, a) = (ba \cdot a)(ab \cdot a) = (ba^2)(ab \cdot a) = (bs)(ab \cdot a)$ or $b(ab \cdot a)$.
5. This follows from Lemma 5.4.
6. $(x, y) = x^{-1}y^{-1} \cdot xy = (x^{-1}y^{-1})(xy^{-1} \cdot y^2) = ((x^{-1}y^{-1})(xy^{-1}) \cdot y^2 = [x^{-2}(xy^{-1}) \cdot (xy^{-1})]y^2 = x^{-2}[(xy^{-1})(xy^{-1})]y^2 = e$ or s . Thus, L is either commutative or else s is its unique non-identity commutator.
 $(x, s) = x^{-1}s^{-1} \cdot xs = s$ implies $x^{-1}s \cdot xs = s$ implies $x^{-1}R_s \cdot xR_s = s$ implies $xJ^2 \cdot x^{-1} = s$ implies $xx^{-1} = s$ implies $s = e$. This is a contradiction, so $(x, s) = e$ implies $s \in C(L, \cdot)$.



Corollary 5.1 *A C -loop with unique non-trivial square is a group.*

Proof

By Lemma 5.4 and Theorem 5.1, L is central square and is of exponent 4. Thus by Theorem 4.3, L is a group. ♠

Remark 5.2 *A C -loop with a unique non-trivial square is an A -loop.*

Theorem 5.2 *Let $(G, \cdot)$ and $(H, \circ)$ be two distinct loop such that the triple $\alpha = (A, B, C)$ is an isotopism of G upon H .*

1. If G is a central square C -loop of exponent 4, then H is a C -loop and an A -loop.
2. If G is a C -loop with a unique non-identity square, then H is a C -loop and an A -loop.

Proof

1. By Theorem 4.3, G is a group and since groups are G-loops, H is a group, thus both a C-loop and an A-loop.
2. The same argument follows by Corollary 5.1.



Remark 5.3 In [18] we considered only isotopes of central loops under triples of the type (A, B, B) and (A, B, A) . But Theorem 5.2 addresses the general triple (A, B, C) where A, B and C are all distinct.

Corollary 5.2 Let $(G, \cdot)$ and $(H, \circ)$ be distinct loops. If the triple (A, B, C) is an isotopism of G upon H such that $(I, L_z^2, JL_z^2J), (R_z^2, I, JR_z^2J) \in \text{AUT}(G, \cdot)$ for all $z \in G$, then H is a C-loop and an A-loop.

Proof

This follows by Theorem 4.2 and Theorem 5.2. ♠

Theorem 5.3 Under a triples of the form (A, A, C) , 'unique non-identity square' is an isotopic invariant property for loops.

Proof

Let $(A, A, C) : (G, \cdot) \rightarrow (H, \circ)$ where G and H are two distinct loops be an isotopism if and only if $xA \circ yA = (x \cdot y)C$. Let $y = x$, $xA \circ xA = (xA)^2 = (x \cdot x)C = x^2C$. If s is the unique non-identity square in G , i.e $x^2 = s$ or $x^2 = e$ for all $x \in G$ then $s' = sC = (xA)^2 = y'^2$ or $y'^2 = (xA)^2 = x^2C = eC = e'$ for all $y' \in H$ with e' as the identity element in H . So, s' is the unique non-identity square element in H . ♠

Corollary 5.3 Central loops with unique non-identity square are isotopic invariant.

Proof

This follows from Theorem 5.2 and Theorem 5.3. ♠

6 Weak and Cross inverse properties in Central Loops

According to [5], the W. I. P. is a generalization of the C. I. P.. The latter was introduced and studied by R. Artzy [3] and [4] while the formal was introduced by J. M. Osborn [22] who also investigated the isotopy invariance of the W. I. P.. Huthnance Jr. [17], proved that the holomorph of a W. I. P. L. is a W. I. P. L.. A loop property is called universal(or at times, a loop is said to be universal relative to a particular property) if the loop has the property and every loop isotope of such a loop possesses such a property. A universal W. I. P. L. is called an Osborn loop. Huthnance Jr. [17] investigated the structure of some holomorph of Osborn loops while Basarab [6] studied Osborn loops that are G-loops. Moufang and conjugacy closed loops have been found to be Osborn loops in [19] but in this section we shall investigate this for C-loops.

Theorem 6.1 *Let $(L, \cdot)$ be an $LC(RC)$ -loop of exponent 3. L is centrum square if and only if L is a C. I. P. L..*

Proof

$x^3 = e$ if and only if $x^2 = x^{-1}$.

$x^2y = yx^2$ if and only if $x^{-1}y = yx^{-1}$ if and only if $x(x^{-1}y) = x(yx^{-1})$ if and only if $y = x(yx^{-1})$ if and only if the C. I. P. holds in the LC-loop L .

$x^2y = yx^2$ if and only if $yx^{-1} = x^{-1}y$ if and only if $(yx^{-1})x = (x^{-1}y)x$ if and only if $y = (x^{-1}y)x$ if and only if the C. I. P. holds in the RC-loop L . ♠

Remark 6.1 *In the proof above, the fact in [4] that in a C. I. P. L., the identity $xy \cdot xJ = y$ and $x(y \cdot xJ) = y$ are equivalent is used. In fact, in a R. I. P. loop L , L has the C. I. P. if and only if $(xJ \cdot y)x = y$.*

Corollary 6.1 *Let $(L, \cdot)$ be an $LC(RC)$ -loop of exponent 3. If L is centrum square, then*

1. L has both the A. I. P. and A. A. I. P.
2. L has the W. I. P.
3. $N = N_\lambda = N_\rho = N_\mu$.
4. $n \in N$ implies $n \in Z(L)$.
5. L is a commutative group.

Proof

1. By Theorem 6.1, L is a C. I. P. L. According to [4] and [5], a C. I. P. L. has the A. I. P. Thus, the first part is true. The second part follows from the fact that $x^2 = x^{-1}$.
2. This follows from the statement in [23] that W. I. P. is a generalization of C. I. P..
- (3) and (4) follow from [5] and [4] hence (5) is as a result of the hypothesis. ♠

Remark 6.2 *In Corollary 6.1, $n \in Z(L)$ for all $n \in N$. Recall that by hypothesis, $n = x^2 \in N$ for all $x \in L$. Thus we expect $x^2 \in Z(L)$ which is actually the hypothesis $x^2 \in C(L)$.*

Corollary 6.2 *Let $(L, \cdot)$ be a C-loop of exponent 3. If L is central square, then*

1. L has the W. I. P. and C. I. P.
2. L is a commutative group.

Proof

A loop $(L, \cdot)$ is a C-loop if and only if it both an LC-loop and an RC-loop. Thence, (1) follows from Theorem 6.1 and Corollary 6.1, while (2) follows from the latter. ♠

Lemma 6.1 *$LC(RC, C)$ -loop of exponent 3 is a group.*

Proof

The proof of this is simple. ♠

Despite the fact that central loops of exponent 3 are groups it will be interesting to study non-commutative central loops of exponent 3 that have the C. I. P. since there exist groups that do not have the C. I. P. From Theorem 6.1, it can be seen that the study of LC(RC)-loops of exponent 3 with C. I. P. is equivalent to the study of centrum square LC(RC)-loops of exponent 3.

The existence of central loops of exponent 3 can be deduced from [15], [26] and [27]. According to [26] and [27], the order of every element in a finite LC(RC)-loop divides the order of the loop. Since $|x| = 3$ for all $x \in L$, $3||L|$ which is possible because:

$$|L| = \begin{cases} 2m, & \text{where } m \geq 3, \quad L \text{ is non-left(right) Bol LC(RC)-loop} \\ 4k, & \text{where } k > 2, \quad L \text{ is non-Moufang but both left(right)-Bol and LC(RC)-loop} \end{cases}$$

The possible orders of finite RC-loops were proved in [27]. In [27], $m > 3$ but in [15], the author gave an example when $m = 1$.

This type of central loops are flexible. This follows from subsection 7.1. RC(LC)-loops are square flexible and $x \mapsto x^{-1}$ is a permutation if and only if $x \mapsto x^2$ is a permutation for a loop of exponent 3. Also, if such finite central loops are extra loops, they will be groups. This is as a result of the claim in [15] that an extra loop in which $x \mapsto x^2$ is a bijection is a group.

Flexible C-loops fall in a class of loops called ARIF loops and has also been studied in [20] and [21]. Authors in [20] constructed a flexible C-loop that is not a RIF loop of order 24 (this is divisible by 3).

6.1 Osborn Central-loops

Theorem 6.2 *Let L be an LC(RC)-loop. L has the R. I. P. (L . I. P.) if and only if has the W. I. P.*

Proof Let $(L, \cdot)$ be a LC-loop with the W. I. P.. Then for all $x, y \in L$, $y(xy)^\rho = x^\rho$. Let $xy = z$, then $x^\lambda(xy) = x^\lambda z$ implies $y = x^\lambda z$, thus $(x^\lambda z)z^\rho = x^\rho$ implies $(x^{-1}z)z^\rho = x^{-1}$. Replacing x^{-1} with x , $(xz)z^\rho = x$. Thence, L has the R. I. P.

Conversely, if L has the I. P., then $y(xy)^\rho = y(xy)^{-1} = y(y^{-1}x^{-1}) = x^{-1} = x^\rho$ hence it has the W. I. P. Let L be a RC-loop with the W. I. P. Then for all $x, y \in L$, $y(xy)^\rho = x^\rho$ if and only if $(xy)^\lambda \cdot x = y^\lambda$. Let $xy = z$, then $(xy)y^\rho = zy^\rho$ implies $x = zy^\rho$. Thus, $z^\lambda(zy^\rho) = y^\lambda$ implies $z^\lambda(zy^{-1}) = y^{-1}$. Replacing y^{-1} by y , $z^\lambda(zy) = y$. Thus, L has the L. I. P..

Conversely, if L has the I. P., then $y(xy)^\rho = y(xy)^{-1} = y(y^{-1}x^{-1}) = x^{-1} = x^\rho$, hence it has the W. I. P. ♠

Corollary 6.3 *Let $(L, \cdot)$ be an LC(RC)-loop with R. I. P. (L . I. P.). Then*

1. $N(L) = N_\lambda(L) = N_\rho(L) = N_\mu(L)$.
2. $(I, R_{x^2}, R_{x^2})((L_{x^2}, I, L_{x^2})) \in \text{AUT}(L)$.

$$3. (L_x^2, R_{x^2}, R_{x^2}L_x^2)((L_{x^2}, R_x^2, L_{x^2}R_x^2) \in AUT(L).$$

Proof

By Theorem 6.2, L has the W. I. P.. According to [22], in a W. I. P. L., the three nuclei coincide, so (1) is true. Thus for an LC-loop, $x^2 \in N_\rho$ and for an RC-loop, $x^2 \in N_\lambda$.

Hence for an LC-loop L , $(L_x^2, I, L_x^2), (I, R_{x^2}, R_{x^2}) \in AUT(L)$ implies $(L_x^2, R_{x^2}, L_x^2 R_{x^2}) = (L_x^2, R_{x^2}, R_{x^2} L_x^2) \in AUT(L)$.

For an RC-loop L , $(I, R_x^2, R_x^2), (L_{x^2}, I, L_{x^2}) \in AUT(L)$ implies $(L_{x^2}, R_x^2, R_x^2 L_{x^2}) = (L_{x^2}, R_x^2, L_{x^2} R_x^2) \in AUT(L)$.

So, (2) and (3) are true. ♠

Remark 6.3 *Corollary 6.3 is true for left(right) Bol loops(i.e LB(RB)-loops). This can be seen from the fact in [23] that a RB(LB)-loop has the L. I. P.(R. I. P.) if and only if it is a Moufang loop which is obviously a W. I. P. L. as mentioned in [19].*

The following facts which is true in Bol loops is also true in central loops.

Theorem 6.3 *A LC(RC)-loop $(L, \cdot)$ is a C-loop if and only if any of the following equivalent statements about L is true.*

1. L has the R. I. P.(L. I. P.)
2. L has the R. A. P.(L. A. P.)
3. L is a RC(LC)-loop.
4. L has the A. A. I. P..(i.e $(xy)^{-1} = y^{-1}x^{-1}$)
5. L has the W. I. P..

Proof

A C-loop satisfies (1)– (2). The proof of the converse is as follows.

1. L is an LC-loop if and only if $(x \cdot xy)z = x(x \cdot yz)$ if and only if $[(x \cdot xy)z]^{-1} = [x(x \cdot yz)]^{-1}$ if and only if $z^{-1}(x \cdot xy)^{-1} = (x \cdot yz)^{-1}x^{-1}$ if and only if $z^{-1}((xy)^{-1} \cdot x^{-1}) = ((yz)^{-1} \cdot x^{-1})x^{-1}$ if and only if $z^{-1}(y^{-1}x^{-1} \cdot x^{-1}) = (z^{-1}y^{-1} \cdot x^{-1})x^{-1}$. Carrying out the replacement x, y, z for x^{-1}, y^{-1}, z^{-1} we have $z(yx \cdot x) = (zy \cdot x)x$ if and only if L is an RC-loop. Hence, L is a C-loop.
2. According to [26], L is an LC-loop if and only if $x \cdot (y \cdot yz) = (x \cdot yy)z$ for all $x, y, z \in L$ while L is an RC-loop if and only if $(zy \cdot y)x = z(yy \cdot x)$ for all $x, y, z \in L$. $x \cdot (y \cdot yz) = (x \cdot yy)z$ if and only if $x \cdot zL_y^2 = xR_{y^2} \cdot z$ if and only if $(R_{y^2}, L_y^{-2}, I) \in AUT(L, \cdot)$ for all $y \in L$ while $(zy \cdot y)x = z(yy \cdot x)$ if and only if $zR^2 \cdot x = z \cdot xL_{y^2}$ if and only if $(R_y^2, L_y^{-1}, I) \in AUT(L, \cdot)$ for all $y \in L$.

If L has the right(left) alternative property, $(R_y^2, L_y^{-2}, I) \in AUT(L, \cdot)$ for all $y \in L$ if and only if L is a C-loop.

3. This is shown in [15].
4. This is equivalent to (1) : If L has the L. I. P.(R. I. P.) then L has the R. I. P.(L. I. P.) implies the A. A. I. P.. Conversely, if L. I. P. holds, then with $z = xy$, we have $y = x^{-1}z$ which gives $z^{-1} = (x^{-1}z)^{-1}x^{-1}$ implies $z^{-1} = (x^{-1}z)^{-1}x^{-1}$ implies $z^{-1} = (z^{-1}x)x^{-1}$. Replacing z^{-1} by z , we have $z = (zx)x^{-1}$ implies the R. I. P.
Similarly, if L has the R. I. P.(L. I. P.) then L has the L. I. P.(R. I. P.) implies the A. A. I. P.. Conversely, if R. I. P. holds, then with $z = xy$, we have $x = zy^{-1}$. Thus, $z^{-1} = y^{-1}(zy^{-1})^{-1} = y^{-1}(yz^{-1})$. Replacing z^{-1} by z , $z = y^{-1}(yz)$ implies L. I. P.
5. This follows from (1) and Theorem 6.2.



Theorem 6.4 ([19]) *The following equivalent conditions define an Osborn loop $(L, \cdot)$.*

1. $x(yz \cdot x) = (x \cdot yE_x) \cdot zx$.
2. $(x \cdot yz)x = xy \cdot (zE_x^{-1} \cdot x)$.
3. $(A_x, R_x, R_xL_x) \in AUT(L)$.
4. $(L_x, B_x, L_xR_x) \in AUT(L)$.

where $A_x = E_xL_x$, $B_x = E_x^{-1}R_x$ and $E_x = R_xL_xR_x^{-1}L_x^{-1}$.

Theorem 6.5 *Let $(L, \cdot)$ be a RC(LC)-loop. If L has the L. I. P.(R. I. P.), then L is an Osborn loop if every element in L is a square.*

Proof

1. Let L be an RC-loop with L. I. P. then by Theorem 6.2, L has the W. I. P..
 $(A_{x^2}, I, L_{x^2}) \in AUT(L)$ if and only if $yA_{x^2} \cdot z = (yz)L_{x^2}$ if and only if $yE_{x^2}L_{x^2} \cdot z = (yz)L_{x^2}$ if and only if $yR_{x^2}L_{x^2}R_{x^2}^{-1}L_{x^2}^{-1}L_{x^2} \cdot z = (yz)L_{x^2}$ if and only if $yR_{x^2}L_{x^2}R_{x^2}^{-1} \cdot z = (yz)L_{x^2}$ if and only if $yR_{x^2}^2L_{x^2}R_{x^2}^{-1} \cdot z = (yz)L_{x^2}$ if and only if $yL_{x^2}R_{x^2}^2R_{x^2}^{-1} \cdot z = (yz)L_{x^2}$ if and only if $yL_{x^2} \cdot z = (yz)L_{x^2}$ if and only if $(L_{x^2}, I, L_{x^2}) \in AUT(L)$ for all $x \in L$ which is true by Corollary 6.3.
Thus, $(I, R_{x^2}^2, R_{x^2}^2)(A_{x^2}, I, L_{x^2}) = (A_{x^2}, R_{x^2}, R_{x^2}L_{x^2}) \in AUT(L)$. Using Theorem 6.4, L is an Osborn loop if every element in L is a square(i.e $y = x^2$ for all $y \in L$).
2. Let L be an LC-loop. If L has the R. I. P., then by Theorem 6.2, L has the W. I. P..
 $(I, B_{x^2}, R_{x^2}) \in AUT(L)$ if and only if $y \cdot zB_{x^2} = (yz)R_{x^2}$ if and only if $y \cdot zE_{x^2}^{-1}R_{x^2} = (yz)R_{x^2}$ if and only if $y \cdot z(R_{x^2}L_{x^2}R_{x^2}^{-1}L_{x^2}^{-1})^{-1}R_{x^2} = (yz)R_{x^2}$ if and only if $y \cdot zL_{x^2}R_{x^2}L_{x^2}^{-1}R_{x^2}^{-1}R_{x^2} = (yz)R_{x^2}$ if and only if $y \cdot zL_{x^2}R_{x^2}L_{x^2}^{-1} = (yz)R_{x^2}$ if and

only if $y \cdot zR_{x^2}L_x^2L_{x^2}^{-1} = (yz)R_{x^2}$ if and only if $y \cdot zR_{x^2} = (yz)R_{x^2}$ if and only if $(I, R_{x^2}, R_{x^2}) \in AUT(L)$ for all $x \in L$ which is true by Corollary 6.3.

Thus, $(L_x^2, I, L_x^2)(I, B_{x^2}, R_{x^2}) = (L_{x^2}, B_{x^2}, L_{x^2}R_{x^2}) \in AUT(L)$. Using Theorem 6.4, L is an Osborn loop if every element in L is a square(i.e $y = x^2$ for all $y \in L$).



Corollary 6.4 *Let $(L, \cdot)$ be a $LC(RC)$ -loop with R . I. $P.(L$. I. $P.)$. L is an Osborn loop if every element in L is a square. Hence, L is a group.*

Proof

This follows from Theorem 6.5. The last conclusion is as a consequence of the fact that $x^2 \in N(L)$. ♠

Corollary 6.5 *Let $(L, \cdot)$ be a C -loop. L is an Osborn loop if every element in L is a square. Hence, L is a group.*

Proof

This follows from Theorem 6.5. The last conclusion is as a consequence of the fact that $x^2 \in N(L)$. ♠

Remark 6.4 *Corollary 6.5 informs us that a class of C -loops that are Osborn are groups. Since the condition attached is not necessary and sufficient, there is the possibility of the existence of C -loops that are Osborn but non-associative.*

7 Question

Does there exist a C -loop that is Osborn but non-associative, non Moufang and non-conjugacy closed.

8 Acknowledgement

The second author would like to express his profound gratitude to the Swedish International Development Cooperation Agency (SIDA) for the support for this research under the framework of the Associateship Scheme of the Abdus Salam International Centre for theoretical Physics, Trieste, Italy.

References

- [1] J. O. Adéníran and A. R. T. Solarin, *A note on automorphic inverse property loops*, Collections of Scientific Papers of the Faculty of Science Krag., **20** 1998, 47–52.
- [2] J. O. Adéníran and A. R. T. Solarin, *A note on generalized Bol identity*, Scientific Annals of Al.I.Cuza. Univ., **45** 1999, 99–102.
- [3] R. Artzy (1955), *On loops with special property*, Proc. Amer. Math. Soc., **6** 1955, 448–453.
- [4] R. Artzy, *Crossed inverse and related loops*, Trans. Amer. Math. Soc., **91(3)** 1959, 480–492.
- [5] R. Artzy, *Relations between loops identities*, Proc. Amer. Math. Soc., **11(6)** 1960, 847–851.
- [6] A. S. Basarab, *Osborn's G-loop*, Quasigroups and Related Systems, **1** 1994, 51–56.
- [7] A. Beg, *A theorem on C-loops*, Kyungpook Math. J. **17(1)** 1977, 91–94.
- [8] A. Beg, *On LC-, RC-, and C-loops*, Kyungpook Math. J. **20(2)** 1980, 211–215.
- [9] R. H. Bruck, *A survey of binary systems*, Springer-Verlag, Berlin-Göttingen-Heidelberg, 1966.
- [10] O. Chein, *A short note on supernuclear (central) elements of inverse property loops*, Arch. Math., **33** 1979, 131–132.
- [11] O. Chein, H. O. Pflugfelder and J. D. H. Smith, *Quasigroups and Loops : Theory and Applications*, Heldermann Verlag, 1990.
- [12] V. O. Chiboka, *Conjugacy Closed Loops of Moufang Type*, Riv. Mat. Pura Appl., **10** 1991, 89–93.
- [13] J. Dene and A. D. Keedwell, *Latin squares and their applications*, the English University press Lts, 1974.
- [14] F. Fenyves, *Extra Loops I*, Publ. Math. Debrecen, **15** 1968, 235–238.
- [15] F. Fenyves, *Extra Loops II*, Publ. Math. Debrecen, **16** 1969, 187–192.
- [16] E. G. Goodaire, E. Jespers and C. P. Milies, *Alternative Loop Rings*, NHMS(184), Elsevier, 1996.
- [17] E. D. Huthnance Jr., *A theory of generalised Moufang loops*, Ph.D. thesis, Georgia Institute of Technology, 1968.
- [18] T. G. Jaiyéolá and J. O. Adéníran : *On Isotopic Characterisation of C-loops*, Pre-Print.

- [19] M. K. Kinyon, *A survey of Osborn loops*, Milehigh conference on loops, quasigroups and non-associative systems, University of Denver, Denver, Colorado, 2005.
- [20] M. K. Kinyon, K. Kunen, J. D. Phillips, *A generalization of Moufang and Steiner loops*, Alg. Univer., **48(1)** 2002, 81–101.
- [21] M. K. Kinyon, J. D. Phillips and P. Vojtěchovský, *C-loops : Extensions and construction*, J. Alg. & its Appl. (to appear).
- [22] J. M. Osborn, *Loops with the weak inverse property*, Pac. J. Math., **10** 1961, 295–304.
- [23] H. O. Pflugfelder, *Quasigroups and Loops : Introduction*, Sigma series in Pure Math. 7, Heldermann Verlag, Berlin, 1990.
- [24] J. D. Phillips and P. Vojtěchovský, *The varieties of loops of Bol-Moufang type*, Alg. Univ., **53(3)** 2005, 115–137.
- [25] J. D. Phillips and P. Vojtěchovský, *The varieties of quasigroups of Bol-Moufang type : An equational reasoning approach* J. Alg., **293** 2005, 17–33
- [26] J. D. Phillips and P. Vojtěchovský, *On C-loops*, Publ. Math. Debrecen, **68(1-2)** 2006, 115–137.
- [27] V. S. Ramamurthi and A. R. T. Solarin, *On finite right central loops*, Publ. Math. Debrecen, **35** 1988, 261–264.
- [28] A. R. T. Solarin, *On the identities of Bol-Moufang type*, Koungpook Math. J., **28(1)** 1998, 51–62.
- [29] A. R. T. Solarin, *On certain Aktivis algebra*, Italian Journal of Pure And Applied Mathematics, **1** 1997, 85–90.
- [30] A. R. T. Solarin and V. O. Chiboka, *A note on G-loops*, Collections of Scientific Papers of the Faculty of Science Krag., **17** 1995, 17–26.

1. Department of Mathematics,
Obafemi Awolowo University,
Ilé Ifè, Nigeria.
e-mail: jaiyeolatemitope@yahoo.com
2. Department of Mathematics,
University of Abeókùta,
Abeókùta 110101, Nigeria.
e-mail: ekenedilichineke@yahoo.com